

# AN EXAMPLE OF A TEICHMÜLLER DISK IN GENUS 4 WITH DEGENERATE KONTSEVICH-ZORICH SPECTRUM

GIOVANNI FORNI AND CARLOS MATHEUS

**ABSTRACT.** We construct an orientable holomorphic quadratic differential on a Riemann surface of genus 4 whose  $SL(2, \mathbb{R})$ -orbit is closed and has a highly degenerate Kontsevich-Zorich spectrum. This example is related to a previous similar construction in genus 3 by the first author.

## 1. INTRODUCTION

The goal of this note is the construction of an orientable quadratic differential  $q$  on a Riemann surface  $M$  of genus 4 such that the Kontsevich-Zorich cocycle along the  $SL(2, \mathbb{R})$ -orbit of  $(M, q)$  has a *totally degenerate* Kontsevich-Zorich spectrum, in the sense that all non-trivial Lyapunov exponents of the Kontsevich-Zorich cocycle are equal to zero. A similar example in genus 3 appeared in [6]. Our example is given by a non-primitive Veech surface in the stratum  $\mathcal{H}(2, 2, 2)$  of abelian differentials with 3 double zeros. Unpublished work of M. Möller implies that there are very few examples of this kind and only for abelian differentials on surfaces of genus  $g \leq 5$ . It is likely that the Forni's example in genus 3 [6] and the example presented here are the only examples of Veech surfaces with vanishing non-trivial Kontsevich-Zorich exponents [11].

In order to explain more precisely our results, let us recall the definition of the Teichmüller flow and the Kontsevich-Zorich cocycle.

Given a surface  $M$ , we denote by  $\text{Diff}^+(M)$  the group of orientation-preserving diffeomorphisms of  $M$  and  $\text{Diff}_0^+(M)$  the connected component of the identity in  $\text{Diff}^+(M)$  (i.e.,  $\text{Diff}_0^+(M)$  is the subset of diffeomorphisms in  $\text{Diff}^+(M)$  which are isotopic to the identity).

**Definition 1.1.** Let  $Q_g$  be the *Teichmüller* space of holomorphic quadratic differentials on a surface of genus  $g \geq 1$ :

$$Q_g = \{\text{holomorphic quadratic differentials}\} / \text{Diff}_0^+(M).$$

Also, let  $\mathcal{M}_g$  be the *moduli* space of holomorphic quadratic differentials on a surface of genus  $g \geq 1$ :

$$\mathcal{M}_g = \{\text{holomorphic quadratic differentials}\} / \Gamma_g,$$

where  $\Gamma_g$  is the *mapping class group*  $\Gamma_g := \text{Diff}^+(M)/\text{Diff}_0^+(M)$ .

Moreover, we note that the group  $SL(2, \mathbb{R})$  acts naturally on  $\mathcal{M}_g$  by linear transformations on the pairs of real-valued 1-forms  $(\text{Re}(q^{1/2}), \text{Im}(q^{1/2}))$ . The *Teichmüller* (geodesic) flow  $G_t$  is given by the action of the diagonal subgroup  $\text{diag}(e^t, e^{-t})$  of  $SL(2, \mathbb{R})$  on  $\mathcal{M}_g$ .

For later reference, we recall some of the main structures of the Teichmüller space  $Q_g$  and the moduli space  $\mathcal{M}_g$ :

- $\mathcal{M}_g$  and  $Q_g$  are stratified into analytic spaces  $\mathcal{M}_\kappa$  and  $Q_\kappa$  obtained by fixing the multiplicities  $\kappa = (k_1, \dots, k_\sigma)$  of the zeroes  $\{p_1, \dots, p_\sigma\}$  of the quadratic differentials (here  $\sum k_i = 4g - 4$ );
- the total area function  $A : \mathcal{M}_g \rightarrow \mathbb{R}^+$ ,  $A(q) = \int_M |q|$  is  $SL(2, \mathbb{R})$ -invariant so that the unit bundle  $\mathcal{M}_g^{(1)} := A^{-1}(1)$  and its strata  $\mathcal{M}_\kappa^{(1)} := \mathcal{M}_g \cap \mathcal{M}_\kappa^{(1)}$  are  $SL(2, \mathbb{R})$ -invariant (and, *a fortiori*,  $G_t$ -invariant);
- a stratum  $\mathcal{M}_\kappa$  of *orientable quadratic differentials* (that is, quadratic differentials obtained as squares of holomorphic 1-forms) has a locally affine structure modeled on the cohomology  $H^1(M, \Sigma_\kappa, \mathbb{C})$ , relative to the zero set  $\Sigma_\kappa := \{p_1, \dots, p_\sigma\}$ , with local charts are given by the period map  $q \mapsto [q^{1/2}] \in H^1(M, \Sigma_\kappa, \mathbb{C})$ ;
- the Lebesgue measure on the Euclidean space  $H^1(M, \Sigma_\kappa, \mathbb{C})$ , appropriately normalized, induces an absolutely continuous  $SL(2, \mathbb{R})$ -invariant measure  $\mu_\kappa$  on  $\mathcal{M}_\kappa$  such that the conditional measure  $\mu_\kappa^{(1)}$  induced on  $\mathcal{M}_\kappa^{(1)}$  is  $SL(2, \mathbb{R})$ -invariant (and hence  $G_t$ -invariant).

Once we get the existence of a good invariant measure  $\mu_\kappa^{(1)}$  for the Teichmüller flow, it is natural to ask whether  $\mu_\kappa^{(1)}$  has finite mass and/or  $\mu_\kappa^{(1)}$  is ergodic with respect to the Teichmüller dynamics. In this direction, Veech [13] showed that the strata are *not* always connected. More recently, Kontsevich and Zorich [8] (in the orientable case) and Laneeau [9] (in the non-orientable case) gave a complete classification of the connected components of all strata of holomorphic quadratic differentials. Taking this into account, we have the following result:

**Theorem 1.2** (Masur [10], Veech [12]). *The total volume of  $\mu_\kappa^{(1)}$  is finite and the Teichmüller flow  $G_t = \text{diag}(e^t, e^{-t})$  is ergodic on each connected component of  $\mathcal{M}_\kappa$  with respect to  $\mu_\kappa^{(1)}$ .*

In order to analyze the Lyapunov spectrum (i.e., the collection of the Lyapunov exponents) of the Teichmüller flow, Kontsevich and Zorich [7] introduced the following notion: the *Kontsevich-Zorich* cocycle  $G_t^{KZ}$  is the quotient of the trivial cocycle  $G_t \times \text{id} : Q_g \times H^1(M, \mathbb{R}) \rightarrow Q_g \times H^1(M, \mathbb{R})$  with respect to the action of the mapping class group  $\Gamma_g$ . It is known that

the cocycle  $G_t^{KZ}$  is *symplectic*, so that the Lyapunov spectrum of  $G_t^{KZ}$  with respect to any  $G_t$ -invariant ergodic probability  $\mu$  is *symmetric*:

$$1 = \lambda_1^\mu \geq \dots \lambda_g^\mu \geq 0 \geq -\lambda_g^\mu \geq \dots \geq -\lambda_1^\mu = -1.$$

It turns out that the  $g$  numbers  $1 = \lambda_1^\mu \geq \lambda_2^\mu \geq \dots \geq \lambda_g^\mu$  appearing in the non-negative part of the Kontsevich-Zorich spectrum determines the Lyapunov spectrum of the Teichmüller flow (this is one of the motivation for introducing the Kontsevich-Zorich cocycle). Indeed, it is possible to show that, for any ergodic probability measure  $\mu$  on  $\mathcal{M}_g$  supported on a stratum of orientable quadratic differentials with  $\sigma \in \mathbb{N}$  distinct zeros, the Lyapunov spectrum of the Teichmüller flow is

$$\begin{aligned} 2 &= (1 + \lambda_1^\mu) \geq \dots \geq (1 + \lambda_g^\mu) \geq \overbrace{1 = \dots = 1}^{\sigma-1} \geq (1 - \lambda_g^\mu) \geq \dots \\ &\geq (1 - \lambda_2^\mu) \geq 0 \geq -(1 - \lambda_2^\mu) \geq \dots \geq -(1 - \lambda_g^\mu) \\ &\geq \underbrace{-1 = \dots = -1}_{\sigma-1} \geq -(1 + \lambda_g^\mu) \geq \dots \geq -(1 + \lambda_1^\mu) = -2. \end{aligned}$$

On the other hand, concerning the Lyapunov spectrum of the Kontsevich-Zorich cocycle, Zorich and Kontsevich conjectured that the Lyapunov exponents of  $G_t^{KZ}$  for the canonical absolutely continuous measure  $\mu_\kappa^{(1)}$  on any stratum of orientable quadratic differentials are all non-zero (i.e., non-uniform hyperbolicity) and distinct (i.e., all Lyapunov exponents have multiplicity 1). After the fundamental works of G. Forni [5] (showing the non-uniform hyperbolicity of  $G_t^{KZ}$ ) and Avila, Viana [1] (proving the simplicity of the Lyapunov spectrum), it follows that Zorich-Kontsevich conjecture is true. In other words, the Lyapunov exponents of a  $\mu_\kappa^{(1)}$ -generic point are all non-zero and they have multiplicity 1. However, it remains to understand the dynamical behavior under  $G_t^{KZ}$  of the non-generic orbits (with respect to  $\mu_\kappa^{(1)}$ ). In particular, one can follow Veech and ask how “degenerate” the Lyapunov spectrum of  $G_t^{KZ}$  can be along a non-typical orbit. This question was first answered by G. Forni [6] who exhibited an example of an orientable holomorphic quadratic differential  $q$  on a Riemann surface  $M$  of genus  $g = 3$  such that the Kontsevich-Zorich spectrum of the  $SL(2, \mathbb{R})$ -invariant measure  $\mu$  supported on the (closed)  $SL(2, \mathbb{R})$ -orbit of  $(M, q)$  verifies  $\lambda_2^\mu = \lambda_3^\mu = 0$ .

At this point, we are able to state our main result:

**Theorem 1.3.** *There exists an orientable holomorphic quadratic differential  $q$  on a Riemann surface  $M$  of genus 4 such that the Lyapunov exponents (with respect to  $G_t^{KZ}$ ) of the  $SL(2, \mathbb{R})$ -invariant probability  $\mu$  supported on*

the  $SL(2, \mathbb{R})$ -orbit of  $(M, q)$  verifies

$$\lambda_2^\mu = \lambda_3^\mu = \lambda_4^\mu = 0.$$

The organization of this note is the following: in §2, we will recall Forni's version of the Kontsevich-Zorich formula for the sum of the Lyapunov exponents of  $G_t^{KZ}$ , in §3 we review Forni's method [6] to exploit symmetries and, in section §4, we will complete the proof of our theorem.

**Acknowledgements:** We would like to thank J.-C. Yoccoz, who motivated our work by asking whether the example in [6] could be generalized. We are also very grateful to M. Möller who introduced us to the notion of a cyclic cover, directed us to the reference [2] and suggested the correct algebraic equation of our example.

## 2. THE KONTSEVICH-ZORICH FORMULA REVISITED

Let  $q$  be a holomorphic quadratic differential on a Riemann surface  $M$  of genus  $g \geq 2$ . Fix  $z = x + iy$  a holomorphic local coordinate and write  $q = \phi(z)dz^2$ . It follows that the (degenerate) Riemannian metric  $R_q$  and the area form  $\omega_q$  induced by  $q$  are  $R_q = |\phi(z)|^{1/2}(dx^2 + dy^2)^{1/2}$ ,  $\omega_q = |\phi(z)|dx \wedge dy$ . Denote by  $S = \partial/\partial x$  and  $T = \partial/\partial y$  the horizontal and vertical directions. Define  $L_q^2(M) := L^2(M, \omega_q)$  the space of complex-valued square-integrable functions and  $H_q^1(M)$  the Sobolev space of functions  $v \in L_q^2(M)$  such that  $Sv, Tv \in L_q^2(M)$ .

**Lemma 2.1** (Forni [4], prop. 3.2). *The Cauchy-Riemann operators  $\partial^\pm := \frac{S \pm iT}{2}$  with (dense) domain  $H_q^1(M) \subset L_q^2(M)$  are closed. Moreover,  $\partial^\pm$  has closed range of finite codimension (equal to the genus  $g$  of  $M$ ). Furthermore, denoting by  $\mathfrak{M}_q^\pm \subset L_q^2(M)$  the subspaces of meromorphic (resp. anti-meromorphic) functions, we have the orthogonal decompositions:*

$$L_q^2(M) = \text{Ran}(\partial_q^+) \oplus \mathfrak{M}_q^- = \text{Ran}(\partial_q^-) \oplus \mathfrak{M}_q^+.$$

Denote by  $\pi_q^\pm : L_q^2(M) \rightarrow \mathfrak{M}_q^\pm$  the orthogonal projection. Let  $H_q$  be the (non-negative definite) Hermitian form on  $\mathfrak{M}_q^+ \subset L_q^2(M)$  given by

$$H_q(m_1^+, m_2^+) := (\pi_q^-(m_1^+), \pi_q^-(m_2^+))_q := \int_M \pi_q^-(m_1^+) \cdot \overline{\pi_q^-(m_2^+)} \omega_q$$

for all  $m_1^+, m_2^+ \in \mathfrak{M}_q^+$ . Let

$$1 \equiv \Lambda_1(q) \geq \Lambda_2(q) \geq \cdots \geq \Lambda_g(q) \geq 0$$

be the eigenvalues of  $H_q$ .

Next, we consider the sets

$$\mathcal{R}_g^{(1)}(k) := \{q \in \mathcal{M}_g^{(1)} : \Lambda_{k+1}(q) = \cdots = \Lambda_g(q) = 0\}.$$

**Definition 2.2.** The set  $\mathcal{R}_g^{(1)}(g-1)$  is called the *determinant locus*.

The relevance of the natural filtration of sets  $\mathcal{R}_g^{(1)}(1) \subset \dots \subset \mathcal{R}_g^{(1)}(g-1)$  for the study of the Lyapunov spectrum of  $G_t^{KZ}$  becomes evident from the following version of a formula by Kontsevich and Zorich [7] for the sum of Lyapunov exponents:

**Theorem 2.3** (Forni [5], Corollary 5.3). *Let  $\mu$  be a  $SL(2, \mathbb{R})$ -invariant ergodic probability measure on  $\mathcal{M}_g^{(1)}$ . Then, the Lyapunov exponents of  $G_t^{KZ}$  with respect to  $\mu$  satisfy the formula:*

$$\lambda_1^\mu + \dots + \lambda_g^\mu = \int_{\mathcal{M}_g^{(1)}} (\Lambda_1(q) + \dots + \Lambda_g(q)) d\mu(q).$$

In particular, since  $\lambda_1^\mu = 1 \equiv \Lambda_1(q)$ , we have

$$\lambda_2^\mu + \dots + \lambda_g^\mu = \int_{\mathcal{M}_g^{(1)}} (\Lambda_2(q) + \dots + \Lambda_g(q)) d\mu(q).$$

A direct consequence of this formula is:

**Corollary 2.4** (Forni [6], Corollary 7.1). *Let  $\mu$  be any  $SL(2, \mathbb{R})$ -invariant ergodic probability measure on  $\mathcal{M}_g^{(1)}$  supported on a stratum of orientable quadratic differentials. The measure  $\mu$  is supported on the locus  $\mathcal{R}_g^{(1)}(1)$  if and only if the non-trivial Kontsevich-Zorich spectrum vanishes, that is,*

$$\lambda_2^\mu = \dots = \lambda_g^\mu = 0.$$

### 3. SYMMETRIES

In this section we recall the simple method developed in [6] to derive bounds on the rank of the matrix  $H_q$  from symmetries of the orientable holomorphic quadratic differential  $q$ . Let  $\text{Aut}(M)$  be the group of holomorphic automorphism of the Riemann surface  $M$ . Let  $\text{Aut}(q) \subset \text{Aut}(M)$  is the subgroup formed by automorphisms  $a \in \text{Aut}(M)$  such that  $a^*(q) = q$ . Note that there is a natural unitary action of  $\text{Aut}(q)$  on  $\mathfrak{M}_q^+$  (by pull-back). Given  $a \in \text{Aut}(q)$ , fix  $\{m_1^+(a), \dots, m_g^+(a)\}$  an orthonormal basis of eigenvectors and denote by  $\{u_1(a), \dots, u_g(a)\}$  the associated eigenvalues. Let  $B^a(q)$  be the matrix of the operator  $\pi_q^- : \mathfrak{M}_q^+ \rightarrow \mathfrak{M}_q^-$  with respect to the basis  $\{m_1^+(a), \dots, m_g^+(a)\} \subset \mathfrak{M}_q^+$  and  $\{\overline{m_1^+(a)}, \dots, \overline{m_g^+(a)}\} \subset \mathfrak{M}_q^-$ :

$$B_{ij}^a(q) = \int_M m_i^+(a) \overline{m_j^+(a)} \omega_q.$$

For any  $I, J \subset \{1, \dots, g\}$  with  $\#I = \#J$ , denote by  $\det B_{IJ}^a$  the minor of the matrix  $B^a(q)$  with entries  $B_{ij}^a(q)$ ,  $i \in I, j \in J$ .

**Lemma 3.1** (Forni [6], Lemma 7.2). *The following holds:*

$$\prod_{i \in I} \prod_{j \in J} u_i(a) u_j(a) \neq 1 \implies \det B_{IJ}^a(q) = 0.$$

**Corollary 3.2.** *If  $\prod_{i \in I} \prod_{j \in J} u_i(a) u_j(a) \neq 1$  for all  $I, J \subset \{1, \dots, g\}$  with  $\#I = \#J = k$ , then  $\text{rank}(H_q) \leq g - k$ , hence  $q \in \mathcal{R}_g^{(1)}(k)$ .*

*Proof.* Since the matrix  $H(q)$  of  $H_q$  with respect to  $\{m_1^+(a), \dots, m_g^+(a)\}$  satisfies  $H(q) = B^a(q)^* B^a(q) = \overline{B^a(q)} B^a(q)$  (see equation (44) of [6]), the statement follows immediately from the previous lemma.  $\square$

At this point, we are ready to present our genus 4 example and prove Theorem 1.3.

#### 4. THE EXAMPLE

Given an integer  $N > 1$  and a 4-tuple of integers  $\mathbf{a} := (a_1, \dots, a_4)$  with  $0 < a_\mu < N$ ,  $\gcd(N, a_1, \dots, a_4) = 1$  and  $\sum_{\mu=1}^4 a_\mu \equiv 0 \pmod{N}$ . Let  $x_1, x_2, x_3, x_4 \in \mathbb{C}$  be 4 distinct points and let  $M := M_N(\mathbf{a})$  be the connected, non-singular Riemann surface determined by the algebraic equation:

$$w^N = (z - x_1)^{a_1} (z - x_2)^{a_2} (z - x_3)^{a_3} (z - x_4)^{a_4}.$$

The surface  $M$  is a *cyclic cover* of the Riemann sphere  $\mathbb{P}^1(\mathbb{C})$  branched over the points  $x_1, \dots, x_4 \in \mathbb{C} \equiv \mathbb{P}^1(\mathbb{C}) \setminus \{\infty\}$ . In fact, every such cyclic cover (up to isomorphisms) can be written as above (see Example 4.2 in [2]). The surface  $M$  has genus

$$g = N + 1 - \frac{1}{2} \sum_{\mu=1}^4 \gcd(a_\mu, N).$$

The surface  $M$  is called a cyclic cover since its automorphism group is cyclic. In fact, it is generated by the automorphism  $T : M \rightarrow M$  given by

$$T(z, w) = (z, \varepsilon w),$$

where  $\varepsilon^N = 1$  is a primitive  $N$ th root of unity.

For  $i \in \{1, \dots, N - 1\}$ , let  $L_i$  denote the eigenspace with eigenvalue  $\varepsilon^i$  for the action of  $T$  on holomorphic abelian differentials:

$$L_i := \{\omega \in H_{\text{dR}}^1(M, \mathbb{C}) : T^* \omega = \varepsilon^i \cdot \omega\}.$$

**Lemma 4.1** (Bouw [2], Lemma 4.3). *The following formulas hold:*

$$\dim_{\mathbb{C}} L_i = \sum_{\mu=1}^4 \left\langle \frac{i \cdot a_{\mu}}{N} \right\rangle - 1$$

(here  $\langle \cdot \rangle$  denotes the fractional part).

We recall that the genus 3 example in [6] was given by the  $SL(2, \mathbb{R})$  orbit of orientable quadratic differentials (or, equivalently, of abelian differentials) constructed from cyclic covers of type  $M_4(1, 1, 1, 1)$ .

*Proof of Theorem 1.3.* Our example is given by an orientable holomorphic quadratic differential constructed from cyclic covers of type  $M_6(1, 1, 1, 3)$ .

Let  $M$  be any branched cover of type  $M_6(1, 1, 1, 3)$ . Observe that  $M$  has genus 4. Lemma 4.1 yields:

$$(4.1) \quad \begin{aligned} \dim_{\mathbb{C}} L_1 &= \dim_{\mathbb{C}} L_2 = 0; \\ \dim_{\mathbb{C}} L_3 &= \dim_{\mathbb{C}} L_4 = 1; \\ \dim_{\mathbb{C}} L_5 &= 2. \end{aligned}$$

Thus, we can fix  $\theta_1 \in L_3$ ,  $\theta_2 \in L_4$  and  $\theta_3, \theta_4 \in L_5$  so that  $\{\theta_1, \dots, \theta_4\}$  is a basis of the space of holomorphic differentials on  $M$  and the action of  $T^*$  on this basis is diagonal with eigenvalues  $-1 = \varepsilon^3$  (with multiplicity 1),  $\varepsilon^4$  (with multiplicity 1) and  $\varepsilon^5$  (with multiplicity 2). Let  $q = \theta_1^2$  be the orientable quadratic differential obtained as pull-back to  $M$  of the unique holomorphic quadratic differential on  $\mathbb{P}(\mathbb{C})$  with simple poles at the branching points  $\{x_1, \dots, x_4\}$ . It follows that the spectrum of the action of  $T \in \text{Aut}(q)$  on the space  $\mathfrak{M}_q^+ \subset L_q^2(M)$  is

$$u_1(T) = 1, \quad u_2(T) = -\varepsilon^4, \quad u_3(T) = -\varepsilon^5, \quad u_4(T) = -\varepsilon^5.$$

Consequently,  $q \in \mathcal{R}_4^{(1)}(1)$ . In fact, a direct application of Corollary 3.2 shows that the form  $H_q$  has rank 1, so that  $q \in \mathcal{R}_4^{(1)}(1)$ .

Let  $\mathcal{V}$  be the set of all orientable quadratic differentials on a cyclic cover of type  $M_6(1, 1, 1, 3)$  branched over 4 points  $\{x_1, \dots, x_4\} \in \mathbb{P}(\mathbb{C})$  obtained as pull-back of the unique (non-orientable) holomorphic quadratic differential on  $\mathbb{P}(\mathbb{C})$ , of total area equal to 1, with simple poles at  $x_1, \dots, x_4$ . Since the stratum of all quadratic differentials on the Riemann sphere with 4 simple poles is  $SL(2, \mathbb{R})$ -invariant and consists of a single  $SL(2, \mathbb{R})$  orbit, the set  $\mathcal{V}$  consists of a single  $SL(2, \mathbb{R})$  orbit in the stratum of orientable quadratic differentials with 3 distinct zeros of order 4 (equivalently, of the stratum of squares of abelian differentials with 3 distinct double zeros). The above computation shows that  $\mathcal{V} \subset \mathcal{R}_4^{(1)}(1)$ , hence (in view of Corollary 2.4) the proof of Theorem 1.3 is completed.  $\square$

We conclude by the remark that our exploration of other cyclic covers has failed to provide additional example of totally degenerate Kontsevich-Zorich spectra. We recall that according to Möller [11], there are no more examples of totally degenerate Kontsevich-Zorich spectrum among Veech surfaces in any genus.

#### REFERENCES

- [1] A. Avila and M. Viana, *Simplicity of Lyapunov Spectra: Proof of the Zorich-Kontsevich conjecture*, Acta Math., v. 146, no. 2, 295–344, 1997.
- [2] I. Bouw, *The  $p$ -rank of ramified covers of curves*, Compositio Math., v. 126, 295–322, 2001.
- [3] I. Bouw and M. Möller, *Teichmüller curves, triangle groups and Lyapunov exponents*, to appear in Ann. of Math.
- [4] G. Forni, *Solutions of the cohomological equation for area-preserving flows on compact surfaces of higher genus*, Ann. of Math., v. 146, no. 2, 295–344, 1997.
- [5] ———, *Deviation of ergodic averages for area-preserving flows on surfaces of higher genus*, Annals of Math., v.155, no. 1, 1–103, 2002.
- [6] ———, *On the Lyapunov exponents of the Kontsevich-Zorich cocycle*, Handbook of Dynamical Systems v. 1B, B. Hasselblatt and A. Katok, eds., Elsevier, 2006, 549–580.
- [7] M. Kontsevich, *Lyapunov exponents and Hodge theory*, in ‘The mathematical beauty of physics’, Saclay, 1996. Adv. Ser. Math. Phys. v. 24, 318–332, World Scientific, River Edge, NJ, 1997.
- [8] M. Kontsevich and A. Zorich, *Connected components of the moduli spaces of Abelian differentials with prescribed singularities*, Invent. Math. v. 153, no. 3, 631–678, 2003.
- [9] E. Lanneau, *Connected components of the strata of the moduli spaces of quadratic differentials*, An. Sc. de l’ENS, v. 41, 1–56, 2008.
- [10] H. Masur, *Interval exchange transformations and measured foliations*, Annals of Mathematics, v. 115, 169–200, 1982.
- [11] M. Möller, personal communication, August 2008.
- [12] W. Veech, *Teichmüller Geodesic Flow*, Annals of Mathematics, v. 124, 441–530, 1986.
- [13] ———, *Moduli spaces of quadratic differentials*, J. Anal. Math., v. 55, 117–171, 1990.

GIOVANNI FORNI: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND,  
COLLEGE PARK, MD 20742-4015, USA  
E-mail address: gforni@math.umd.edu.

CARLOS MATHEUS: COLLÈGE DE FRANCE, 3, RUE D’ULM, PARIS, CEDEX 05,  
FRANCE  
E-mail address: matheus@impa.br.